

University of Groningen

Molecular aspects of antibody-antigen interactions

Schellekens, Gerardus Antonius

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

1995

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Schellekens, G. A. (1995). *Molecular aspects of antibody-antigen interactions: size reduction of a herpes simplex virus neutralizing antibody and its antigen*. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Summary

Summary

Antibody molecules, produced as a response against foreign substances, interact with their antigen in a very specific manner. Antibodies with a predetermined specificity (monoclonal antibodies) can be produced and are widely used in medicine and science as indicator molecules. Genetic engineering of antibody molecules offers the possibility to study the nature of the antibody-antigen interaction in more detail and to modify antibody molecules for specific applications.

Only a small part of the antibody molecule is directly involved in antigen binding. This antigen binding site is made up of the variable domains of the heavy- and light chain. Also, only part of the antigen molecule is involved in the antibody-antigen interaction. The part of the antigen that is involved in the interaction with the antibody is referred to as the 'epitope'. For some applications size reduction of the antibody molecule could be beneficial, e.g. a more stable ligand in biosensor applications and in immunoaffinity chromatography, and for structural investigations with NMR spectroscopy.

In this thesis size reduction of both the antibody and the antigen are studied in order to identify the minimal parts of the molecules involved in the interaction. For this purpose the monoclonal antibody A16, that recognizes an epitope on glycoprotein D of herpes simplex virus, was used as a model system. Several approaches for size reduction of the A16 antibody were studied simultaneously. Antibody fragments of different size derived from the variable domains, either produced in *E. coli* or as synthetic peptides, were evaluated. The soluble expression of recombinant fragments proved to be very difficult. Unfortunately, neither the recombinant fragments nor the synthetic peptides showed a detectable affinity towards the antigen. Hence, antigen binding of fragments smaller than the Fv (made up of the variable domains of heavy- and light chain) is only encountered in very special cases.

In order to identify the minimal part of the antigen involved in the interaction with the antibody, A16 was used to select binding peptides from a large library of random peptides that are displayed on the surface of a filamentous bacteriophage. A binding pattern of three essential amino acid residues was identified. With the use of synthetic peptides and biosensor technology a seven residue peptide was identified as the minimal antibody binding site. A peptide derived from the library displayed similar binding- and immunological characteristics as a peptide derived from the antigen (glycoprotein D). This method failed to select binding peptides

using the non-antibody target myoglobin. During these studies a method was developed to study the binding characteristics of selected phage clones, prior to DNA sequencing and peptide synthesis, using biosensor technology.